
ON EXISTENCE OF WEAKLY MAXIMAL PROGRAMS IN VON NEUMANN GROWTH MODELS

by

WILLIAM A. BROCK

AD 670486



OPERATIONS RESEARCH **CENTER**

This document has been approved for public release and sale; its distribution is unlimited

UNIVERSITY OF CALIFORNIA . BERKELEY

ON EXISTENCE OF WEAKLY MAXIMAL PROGRAMS IN VON NEUMANN GROWTH MODELS

by

William A. Brock
Operations Research Center
University of California, Berkeley

April 1968 ORC 68-10

This research has been partially supported by the Office of Naval Research under Contract Nonr-222(83), and the National Science Foundation under Grant GP-7417 with the University of California. Reproduction in whole or in part is permitted for any purpose of the United States Government.

ACKNOWLEDGEMENT

The author wishes to thank Dr. David Gale for valuable assistance and encouragement.

Dr. Gale suggested investigating whether the optimal stationary program was optimal in the class of programs starting from it. This turned out not to be true but the investigation of it lead to the present work.

ABSTRACT

We consider, with D. Gale, an n-sector economy defined by a convex, compact set T consisting of input-output pairs $(x,y) \in \mathbb{R}^{2m}$ where \mathbb{R}^{2m} is real 2m-dimensional space. On T is defined a concave, continuous real function u that associates utility to each input-output pair. T and u are called the technology and the utility function of the economy. At each $(x,y) \in T$ we assume u has bounded steepness, i.e.,

 $\sup_{(x',y')\in T} | u(x',y') - u(x,y) | / | | (x',y') - (x,y) | | < \infty$

where $||\cdot||$ is any norm on R^{2m} . A program with initial stocks y_0 is a sequence $\left\{(x_n,y_n)\right\}_{n=1}^{\infty}$ of points of T such that $x_n \leq y_{n-1}$, $n=1,2,\ldots$ An optimal stationary program (OSP) is a pair $(\bar{x},\bar{y}) \in T$ such that (\bar{x},\bar{y}) solves: $\max u(x,y)$ over $\{(x,y) \in T \mid x \leq y\}$ where \leq is componentwise ordering on R^m . The initial stock y_0 is sufficient if there is a program $\left\{(x_n,y_n)\right\}_{n=1}^{\infty}$ from y_0 such that $y_N > 0$ for some N. The technology T is productive if there is $(x,y) \in T$ such that x < y (i.e., $x_1 < y_1$, $i = 1,2,\ldots,m$). We say inaction is possible if $(0,0) \in T$. A program $\{(x_n',y_n')\}$ from y_0 is optimal (weakly maximal) if

$$\frac{\lim_{N\to\infty}\sum_{n=1}^{N} (u(x_n',y_n') - u(x_n,y_n)) \ge 0}{0\left(\frac{\lim_{N\to\infty}\sum_{n=1}^{N} (u(x_n',y_n') - u(x_n,y_n)) \ge 0\right)}$$

for all programs $\{(x_n,y_n)\}$ starting from y_0 . The utility function u is strictly concave at $(\bar{x},\bar{y}) \in T$ if $u(\lambda(\bar{x},\bar{y}) + (1-\lambda)(x,y)) > \lambda u(\bar{x},\bar{y}) + (1-\lambda)u(x,y)$ for all $(x,y) \neq (\bar{x},\bar{y})$, $(x,y) \in T$, $0 < \lambda < 1$.

An easy proof is given of Gale's result: If y_0 is sufficient, T productive, inaction is possible, and u is strictly concave at (\bar{x},\bar{y}) then there is an optimal program from y_0 . The main result is: If (\bar{x},\bar{y}) is the unique OSP, y_0 is sufficient, T is productive, and inaction is possible then there is a weakly maximal program from y_0 . An example is given where the hypotheses of the main result are satisfied but the OSP does not outgrow every program starting from it.

The importance of the main result lies in removing the objectionable hypothesis of strict concavity on the utility function.

ON EXISTENCE OF WEAKLY MAXIMAL PROGRAMS IN VON NEUMANN GROWTH MODELS

by

William A. Brock

In this paper we shall consider general models of economic growth, formulate an "optimality" criterion, and show that there exist "optimal programs" under weak, economically meaningful assumptions. We now turn to development of the model following Gale [1].

Let T be a compact, convex set in real 2m space R^{2m} ($R^{2m} \equiv \{(x,y) \mid x \text{ is a real m-vector and } y \text{ is a real m-vector}\}$). Let u be a concave, continuous numerical function defined on T. T represents the possible input-output combinations and u attaches utility to each. T,u are called the technology and the utility function respectively. Assume further that u has bounded steepness over T, i.e., sup $\sigma(x,y) < \infty$ where $(x,y) \in T$

$$\sigma(x,y) \equiv \sup_{(x',y') \in T} |u(x',y') - u(x,y)|/||(x',y') - (x,y)||$$

for each $(x,y) \in T$, where $||\cdot||$ is a norm on R^{2m} . Bounded steepness at (x,y), i.e., $\sigma(x,y) < \infty$ means that no infinitesimal movement from (x,y) can produce a large change in utility.

Let $0 \le y_0 \in \mathbb{R}^m$. A program with initial stocks y_0 is a sequence $\{(x_n,y_n)\}_{n=1}^\infty$ of points $(x_n,y_n)\in T$ such that $x_n \le y_{n-1}$, $n=1,2,\ldots$ With each program there is associated a utility sequence $\{u(x_n,y_n)\}_{n=1}^\infty$.

The problem of optimal economic growth is: given initial stocks y_0 , find a "reasonable" partial ordering of utility sequences generated by the economy and show there exist maximal elements with respect to this ordering.

It is an easy exercise to show $V(x,y) \in T$, $\sigma(x,y) < \infty$ implies u(x,y) has bounded steepness over T.

We say that $\{(x_n,y_n)\}$ overtakes (strictly overtakes) $\{(x_n^i,y_n^i)\}$ if

$$\frac{\lim_{N\to\infty}\left(\sum_{n=1}^{N}u(x_{n},y_{n})-u(x_{n}',y_{n}')\right)\geq0, (>0)$$
 (1)

 $\{(\mathbf{x_n}, \mathbf{y_n})\}$ is optimal if it overtakes every other program starting from $\mathbf{y_o}$. $\{(\mathbf{x_n}, \mathbf{y_n})\}$ is maximal (weakly maximal) if no other program starting from $\mathbf{y_o}$ overtakes (strictly overtakes) it. Overtaking is clearly a partial ordering on the set of programs starting from $\mathbf{y_o}$ and a maximal program is a maximal element with respect to this ordering. We now introduce the notion of optimal stationary program (OSP) to facilitate construction of weakly maximal programs.

Definition:

 $(\bar{x},\bar{y}) \in T$ is an OSP if (\bar{x},\bar{y}) solves:

$$\max \{u(x,y) \mid x \leq y, (x,y) \in T\}.$$

Let us assume,

Assumption 1:

T is productive and inaction is possible, i.e., there $(x,y) \in T$ such that x < y, and $(0,0) \in T$, respectively.

Using productivity of T we may prove

Lemma 1

If (\bar{x},\bar{y}) is an OSP and T is productive then there is $p \ge 0$, $p \in R^m$ such that (\bar{x},\bar{y}) , p solve:

Maximize
$$u(x,y) + p \cdot (y - x)$$

Subject to $(x,y) \in T$

furthermore

$$p \cdot (\bar{y} - \bar{x}) = 0.$$

Proof:

This is an application of the Kuhn-Tucker Theorem to the problem of maximizing u(x,y) subject to the constraint $x \le y$ where, by productivity of T, the constraint can be satisfied strictly and this is sufficient (see [3]) for the validity of the Kuhn-Tucker Theorem.

Let us now consider an important example, i.e., the "Von Neumann" economy:

$$u(x,y) \equiv max \{c.v \mid Av = x, Bv = y\}, T \equiv \{Av,Bv\} \mid v \in K\}$$

where A,B are m x k nonnegative goods matrices and $K \subseteq \mathbb{R}^k$ is a compact polyhedron (e.g., labor limitations) constraining activity levels v and $c \cdot v = \sum_{i=1}^k c_i v_i$ is the "value" or utility obtained from operating the k activities at intensity v.

We want to find reasonable assumptions so that this economy has maximal or weakly maximal programs. In [1] Gale shows that strict concavity of u at (\bar{x},\bar{y}) implies existence of an optimal program from y_0 . But the utility function of the Von Neumann economy is strictly concave at no $(x,y) \in T$.

If one is satisfied with a weakly maximal program then it is sufficient to assume that the OSP, (\bar{x},\bar{y}) , is unique and to make some mild assumptions on T, y_0 . We shall now work toward the proof of this result.

Assumption 2:

 y_o is sufficient, i.e., there exists a program $\{(x_n,y_n)\}_{n=1}^\infty$ from y_o such that † $y_n>0$ for some n .

[†]Given a vector $x \in \mathbb{R}^m$, x > 0 ($x \ge 0$) means $x_i > 0$ ($x_i \ge 0$) i = 1, 2, ..., m.

We now define an important class of programs.

Definition:

A program is good if there exists a real number G such that

$$G \leq \sum_{n=1}^{N} u(x_n, y_n) - u(\bar{x}, \bar{y}) ,$$

for all positive integers N .

For the sake of completeness we state and give Gale's proof of the existence of good programs.

Lemma 2 (Gale)

If y_0 , T satisfy Assumptions 1,2, then there is a good program from y_0 .

Proof:

Since y_0 is sufficient we can, after finite time, produce a bundle $y_n>0$. Since goodness does not depend on what happens in a finite amount of time we may as well assume $y_0>0$. Using $(0,0)\in T$ and convexity of T choose a productive pair $(x_p,y_p)\in T$ such that $x_p< y_0$. Choose $0\le \lambda<1$ such that $x_1\equiv (1-\lambda)|x+\lambda x_p\le y_0$, $y_1\equiv (1-\lambda)|y+\lambda y_p$. Define the sequence $\{(x_n,y_n)\}$ by

$$(x_n,y_n) = (1 - \lambda^n)(\bar{x},\bar{y}) + \lambda^n(x_p,y_p)$$
, n = 1, 2, ...

Using the recursion:

$$(x_n,y_n) = (1 - \lambda)(x,y) + \lambda(x_{n-1},y_{n-1}), n = 2, 3, ...$$

it is easy to check that $\{(x_n,y_n)\}$ is a program from y_0 . Using bounded steepness we get, for all N ,

$$\begin{split} \sum_{n=1}^{N} |u(\bar{x}, \bar{y}) - u(x_n, y_n)| &\leq \sum_{n=1}^{N} |\sigma(x_n, y_n)| |(\bar{x}, \bar{y}) - (x_n, y_n)| |\\ &\leq \sum_{n=1}^{N} |\sup_{(x, y) \in T} |\sigma(x, y)| |\lambda^n| ||(x_p - \bar{x}, y_p - \bar{y})| |\\ &< \infty \text{, since } \sup_{(x, y) \in T} |\sigma(x, y)| < \infty \text{, } 0 \leq \lambda < 1 \text{.} \end{split}$$

It follows that $\{(x_n,y_n)\}$ is good. It turns out that

Lemma 3 (Gale)

A program is good or its associated series converges to $-\infty$.

Proof:

From this point on we set $u(\bar{x},\bar{y})=0$ to ease the writing. If a program is not good then for all real numbers G there is M depending on G such that

$$\sum_{n=1}^{M} u(x_n, y_n) \leq G.$$

By Lemma 1 and compactness of T

$$\sum_{n=M+1}^{N} u(x_n, y_n) \le \sum_{n=M+1}^{N} p \cdot (x_n - y_n) \le p \cdot (x_{M+1} - y_N) < B$$

where B is a bound independent of M and N . Hence

$$\sum_{n=1}^{N} u(x_n, y_n) \le G - B \qquad \text{for } N \ge M.$$

It follows that

$$\sum_{n=1}^{N} u(x_n, y_n) \rightarrow -\infty , N \rightarrow \infty .$$

"Turnpike" properties appear to be everpresent in the theory of optimal economic growth. We prove an "average turnpike" property for good programs.

Lemma 4

If $\{(\mathbf{x}_n,\mathbf{y}_n)\}$ is good then $\mathbf{u}(\overline{\mathbf{x}}_n,\overline{\mathbf{y}}_n) + \mathbf{u}(\overline{\mathbf{x}},\overline{\mathbf{y}})$ where $(\overline{\mathbf{x}}_n,\overline{\mathbf{y}}_n) \equiv \frac{1}{n} \sum_{i=1}^n (\mathbf{x}_i,\mathbf{y}_i) \in \mathbf{T} . \text{ In particular, if } (\overline{\mathbf{x}},\overline{\mathbf{y}}) \text{ is the unique OSP then } (\overline{\mathbf{x}}_n,\overline{\mathbf{y}}_n) + (\overline{\mathbf{x}},\overline{\mathbf{y}}) .$

Proof:

By goodness of $\{(x_n,y_n)\}$ and concavity of u , there is G such that for each N

$$\frac{G}{N} \leq \frac{1}{N} \sum_{n=1}^{N} u(x_n, y_n) \leq u(\bar{x}_N, \bar{y}_N).$$

Let (x,y) be any cluster point of the sequence $\{(x_n,y_n)\}$. Then $0 \le u(x,y)$, $x \le y$, and $(x,y) \in T$ because

$$(\bar{x}_n, \bar{y}_n) = \frac{1}{n} \sum_{i=1}^{n} (x_i, y_i) \in T$$
, $n = 1, 2, ...$

and T is closed. Hence (x,y) is an OSP and

$$u(\bar{x}_n, \bar{y}_n) \rightarrow u(\bar{x}, \bar{y}) \equiv 0 , n \rightarrow \infty$$
.

In particular if (\bar{x},\bar{y}) is unique, $(x,y) = (\bar{x},\bar{y})$, thus

$$(\bar{x}_n,\bar{y}_n) \rightarrow (\bar{x},\bar{y})$$
 , $n \rightarrow \infty$.

After proving one more Lemma we may prove the main result of this paper.

Lemma 5

If y_0 , T satisfy Assumptions 1,2 then there is a nonnegative sequence $\{\delta_n\}_{n=1}^{\infty}$ associated with $\{u(x_n,y_n)\}$ such that

$$\sum_{n=1}^{N} u(x_n, y_n) = p \cdot (y_0 - y_N) - \sum_{n=1}^{N} \delta_n, N = 1, 2, ...$$
 (1)

Furthermore, a program $\{(x_n',y_n')\}$ may be found so that its associated series $\sum_{n=1}^\infty \delta_n' \quad \text{is minimal in the class of programs starting from } y_0 \ .$

Proof:

By Lemma 1 there is $\beta_n \ge 0$ such that

$$u(x_n, y_n) = p \cdot (x_n - y_n) - \beta_n, n = 1, 2, ...$$
 (2)

Summing (2) to N and setting $\delta_n = -p \cdot (x_n - y_{n-1}) + \beta_n \ge 0$ gives (1). For $\{(x_n, y_n)\}$ good there exist G, B such that

$$G \le \sum_{n=1}^{N} u(x_n, y_n) = p \cdot (y_0 - y_N) - \sum_{n=1}^{N} \delta_n \le B - \sum_{n=1}^{N} \delta_n, N = 1, 2, ...$$

Thus

$$\sum_{n=1}^{N} \delta_{n} \leq B - G , N = 1, 2, ...$$

so that $\sum\limits_{n=1}^\infty \delta_n < \infty$ for good programs. Our task is to find $\{(x_n',y_n')\}$ starting from y_o such that $\alpha = \inf \left\{\sum\limits_{n=1}^\infty \delta_n : \{\delta_n\} \text{ is associated with } \{(x_n,y_n)\} \text{ starting from } y_o\right\} = \sum\limits_{n=1}^\infty \delta_n'$. By existence of good programs (Lemma 2) we have $\alpha < \infty$.

Choose a "minimizing" sequence of programs $\{(x_n^N, y_n^N)\}$ such that

$$\sum_{n=1}^{\infty} \delta_n^{N} \leq \alpha + \frac{1}{N}, N = 1, 2, \dots$$

By compactness of T , use the Cantor diagonal process to select a subsequence $\{N^*\} \subseteq \{N\}$ such that for each in

$$\left(x_n^{N'}, y_n^{N'}\right) \rightarrow \left(x_n^{i}, y_n^{i}\right)$$
, $N' \rightarrow \infty$.

It is easy to check that $\{(x_n',y_n')\}$ is a program. By compactness of T and continuity of u, $\{\delta \frac{N'}{n}\}_{N'}$ is bounded for each n. Hence we may use the Cantor diagonal process again to choose a subsequence $\{N''\} \subseteq \{N'\}$ so that

$$\delta_n^{N''} \rightarrow \delta_n'$$
 , $n = 1, 2, \ldots$

It is clear from continuity of u and definition of δ_n that $\{\delta_n^*\}$ corresponds to the program $\{(x_n^*,y_n^*)\}$. Hence

$$\sum_{n=1}^{\infty} \delta_n^* \geq \alpha.$$

Suppose $\beta \equiv \sum_{n=1}^{\infty} \delta_n' > \alpha$. Let r_1 , r_2 be chosen so that $\alpha < r_1 < r_2 < \beta$. Choose N_0 so that $N \geq N_0$ implies $\sum_{n=1}^{\infty} \delta_n' \geq r_2$. Choose M_0 (depending on N_0)

so that $M \in \{N''\}$, $M \ge M_0$ implies $\sum_{n=1}^N \delta_n^M \ge r_1$. But $\alpha + \frac{1}{M} \ge \sum_{n=1}^\infty \delta_n^M \ge \sum_{n=1}^\infty \delta_n^M$. Hence there are infinitely many $M \in \{N''\}$ such that $\alpha + \frac{1}{M} \ge r_1 > \alpha$. This is a contradiction. Thus

$$\alpha = \sum_{n=1}^{\infty} \delta_n^{\dagger} .$$

We now prove our main result

Theorem 1

If there exists a unique OSP and T, y_0 satisfy Assumptions 1,2 then there is a weakly maximal program from y_0 . If $y_0 = \bar{y}$ then a weakly maximal program is (\bar{x}, \bar{y}) .

Proof:

Let $\{(x_n',y_n')\}$ be a program with minimal $\sum\limits_{n=1}^\infty \delta_n'$. From (1) of Lemma 5 we get

$$\sum_{n=1}^{N} u(x_{n}, y_{n}) - u(x_{n}', y_{n}') = p \cdot (y_{N}' - y_{N}) + \sum_{n=1}^{N} \delta_{n}' - \sum_{n=1}^{N} \delta_{n}, N = 1, 2, ...$$
 (1)

for any other program $\{(x_n,y_n)\}$. Assume $\{(x_n,y_n)\}$ strictly outgrows $\{(x_n',y_n')\}$. Using (1) above there are M , $\lambda>0$ such that

$$\lambda \leq p \cdot (y_N' - y_N), N \geq M$$
 (2)

Since we need only compare good programs (Lemma 2) we may use Lemma 4 (the average turnpike Lemma) to obtain

$$\lambda \leq \frac{1 \text{im}}{N + \infty} p \cdot (\bar{y}_N^* - \bar{y}_N^*) = 0$$

which is a contradiction. If $\bar{y} = y_0$ then $\sum_{n=1}^{\infty} \delta_n' = 0$ if $\{\delta_n'\}$ is associated with $\{(\bar{x},\bar{y})\}$. Hence (\bar{x},\bar{y}) is weakly maximal (note that, if $y_0 = \bar{y}$, the assumption that y_0 is sufficient is not needed for the weak maximality proof).

One might ask if it is possible to strengthen Theorem 1 by showing existence of a strongly optimal program, i.e.,

$$\frac{\lim_{N\to\infty} \sum_{n=1}^{N} u(x_n, y_n) - u(x_n', y_n') > 0 \qquad \text{for all } \{(x_n', y_n')\}.$$

We produce a counterexample where the OSP is unique, but there is no strongly optimal program from \bar{y} .

We consider a one good, two activity Von Neumann economy. Let $v_1 \ge 0$, $v_2 \ge 0$ be activity levels of A_1 , A_2 where A_1 doubles capital but produces no utility, A_2 halves capital and produces one unit of utility. Let the labor constraint be $v_1 + v_2 \le 1$. Then A = (1,2), B = (2,1). Solving the problem:

Maximize
$$v_1 + v_2 \le v_1 + v_2$$

Subject to $v_1 + 2v_2 \le 2v_1 + v_2$; $v_1 + v_2 \le 1$, $v_1 \ge 0$, $v_2 \ge 0$

gives us the unique OSP

$$\vec{v}_1 = \vec{v}_2 = \frac{1}{2}$$
, $\vec{x} = \vec{v}_1 + 2\vec{v}_2 = \frac{3}{2}$, $\vec{y} = 2\vec{v}_1 + \vec{v}_2 = \frac{3}{2}$, $\vec{u} = \vec{v}_2 = \frac{1}{2}$.

Starting from \bar{y} we have the program: (to see that the following is a program check that $Av_n \leq Bv_{n-1}$, $n=1, 2, \ldots$ where $v_o = \bar{v}$)

$$(v_1^n, v_2^n) = (\frac{1}{6}, \frac{2}{3}); u_1 = \frac{2}{3}$$

 $(v_1^n, v_2^n) = (1, 0), n \text{ even } ; u_n = 0$
 $(v_1^n, v_2^n) = (0, 1), n \text{ odd } ; u_n = 1, n > 1$

Note that

$$\overline{\lim} \sum_{n=1}^{N} (u_n - \overline{u}) > 0$$

so that (\bar{x},\bar{y}) does not outgrow $\{(x_n,y_n)\}$. However, by Theorem 1 (\bar{x},\bar{y}) is weakly maximal. Hence there can be no strongly optimal program from (\bar{x},\bar{y}) . To get optimal programs it is sufficient to assume strict concavity of v at (\bar{x},\bar{y}) .

The strong result of Gale may be obtained easily by the method of proof employed in this paper.

Theorem 2 (Gale)

If u is strictly concave at (\bar{x},\bar{y}) and y_0 , T satisfy Assumptions 1,2 then there is a program $\{(x_n',y_n')\}$ from y_0 such that for each $\{(x_n,y_n)\}$ starting from y_0

$$\frac{\lim_{N\to\infty}\sum_{n=1}^{N}u(x_{n}',y_{n}')-u(x_{n},y_{n})\geq 0$$
 (1)

Proof:

Select $\{(x_n',y_n')\}$ with minimal $\sum\limits_{n=1}^\infty \delta_n'$. It is easy to check that $\{(x_n',y_n')\}$ is good. By Lemma 3 we need only compare good programs. Consider

[†]A function f on a set X is strictly concave at $\bar{x} \in X$ if for each $x \in X$, $0 < \lambda < 1$, $f(\lambda \bar{x} + (1 - \lambda)x) > \lambda f(\bar{x}) + (1 - \lambda)f(x)$.

a good program $\{(x_n,y_n)\}$. We show $(x_n,y_n) \rightarrow (\overline{x},\overline{y})$, $n \rightarrow \infty$. Since

$$\sum_{n=1}^{N} u(x_{n}, y_{n}) = \sum_{n=1}^{N} p \cdot (x_{n} - y_{n}) - \sum_{n=1}^{N} \beta_{n} \leq p \cdot (y_{0} - y_{N}) - \sum_{n=1}^{N} \beta_{n}$$

where $0 \le \beta_n = p(x_n - y_n) - u(x_n, y_n)$, there is G (since $\{(x_n, y_n)\}$ is good) such that

$$\sum_{n=1}^{N} \beta_{n} \le p \cdot (y_{0} - y_{N}) - G \le B - G, \qquad N = 1, 2, ...$$

Therefore $\beta_n \to 0$. Now let (\bar{x},\bar{y}) be any cluster point of $\{(x_n,y_n)\}$. Since $\beta_n \to 0$, therefore $u(\bar{x},\bar{y})+p\cdot(\bar{y}-\bar{x})=0$. But strict concavity of u at (\bar{x},\bar{y}) implies (\bar{x},\bar{y}) is the unique maximizer over T of the nonpositive function, $u(x,y)+p\cdot(y-x)$. Thus $(\bar{x},\bar{y})=(\bar{x},\bar{y})$ and $(x_n,y_n)+(\bar{x},\bar{y})$ as $n\to\infty$. Now, as in Theorem 1, let $\{(x_n',y_n')\}$ be a program with minimum $\sum_{n=1}^\infty \delta_n'$. Since $y_n'+\bar{y}$, $y_n\to \bar{y}$ for good programs it is now clear from (1) of Theorem 1 that

$$\frac{1im}{N\to\infty}\sum_{n=1}^{N}u(x_n',y_n')-u(x_n,y_n)\geq 0.$$

One might expect weakly maximal programs to exist in almost any model.

Gale's [1] "cake eating" example with u strictly concave has no weakly maximal program. McKenzie [2] constructed an example of a Von Neumann model with linear utility that has no weakly maximal programs. Both of these examples have multiple OSP's. We doubt that it is possible to replace "weakly maximal" in Theorem 1 by "maximal".

BIBLIOGRAPHY

- [1] Gale, D., "On Optimal Development in a Multisector Economy," Review of Economic Studies, Vol. 34, pp. 1-18, (1967).
- [2] Private Communication of L. McKenzie to D. Gale.
- [3] Karlin, S., MATHEMATICAL METHODS AND THEORY IN GAMES, PROGRAMMING AND ECONOMICS, Vol. I, Addison-Wesley, Reading, Massachusetts, pp. 200-201, (1959).

Security Classification

DOCUMENT CO (Security classification of title, body of abstract and index	NTROL DATA - R&		the everall report is clausified)	
1. ORIGINATING ACTIVITY (Corporate author)	me amoration against of ar		RI SECURITY CLASSIFICATION	
University of California, Berkeley		Unc	lassified	
oniversity of California, Derkeley		P		
3. REPORT TITLE		<u> </u>	· · · · · · · · · · · · · · · · · · ·	
ON EXISTENCE OF WEAKLY MAXIMAL PROGRA	MS IN VON NEUMA	INN GRO	WTH MODELS	
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)				
Research Report S. AUTHOR(S) (Lest name. liret name, initial)				
BROCK, William A.				
	•			
6. REPORT DATE	76. TOTAL NO. OF P	AGES	76. NO. OF REFS	
April 1968	13		3	
Nonr-222(83)	90 URIGINATOR'S RE			
b. PROJECT NO.	ORC 68-10			
NR 047 033				
с.	Sb. OTHER REPORT	NO(\$) (Any	other numbers that may be assigned	
Research Project No.: RR 003 07 01				
10 A VAIL ABILITY/LIMITATION NOTICES				
Distribution of this document is unlin	mited.			
11. SUPPLEMENTARY NOTES Also supported by	12. SPONSORING MILIT	TARY ACTI	IVITY	
the National Science Foundation under				
Grant GP-7417.	Mathema	tical S	Science Division	
	<u> </u>			
13 ABSTRACT				
SEE ABSTRACT.				
			·	

DD .5284. 1473

Unclassified
Security Classification

KEY WORDS	LINK A		LINK B		LINKC	
	MOLE	WT	ROLE	WT	ROLE	WT
Dynamic Programming						
Economic Planning with an Infinite Time Horizon						

INSTRUCTIONS

- 1. ORIGINATING ACTIVITY: Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (corporate author) issuing the report.
- 2a. REPORT SECURITY CLASSIFICATION: Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.
- 2b. GROUP: Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.
- 3. REFORT TITLE: Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.
- 4. DESCRIPTIVE NOTES: If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.
- 5. AUTHOR(S): Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.
- 6. REPORT DATE: Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.
- 7a. TOTAL NUMBER OF PAGES: The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.
- 7b. NUMBER OF REFERENCES: Enter the total number of references cited in the report.
- 8a. CONTRACT OR GRANT NUMBER: If appropriate, enter the applicable number of the contract or grant under which the report was written.
- 8b, 8c, & 8d. PROJECT NUMBER: Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.
- 9a. ORIGINATOR'S REPORT NUMBER(S): Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.
- 9b. OTHER REPORT NUMBER(S): If the report has been assigned any other report numbers (either by the originator or by the sponsor), also enter this number(s).
- 10. AVAILABILITY/LIMITATION NOTICES: Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known

- 11. SUPPLEMENTARY NOTES: Use for additional explanatory notes.
- 12. SPONSORING MILITARY ACTIVITY: Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address.
- 13. ABSTRACT: Enter an abstract giving a brief and factual numbers of the document indicative of the report, even though it may also appear alsowhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS). (S). (C), or (U)

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. KEY WORDS: Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classifulation in required. Identifiers, such as equipment model designation, tride name, military project code name, genyraphic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.